

Dark matter, the CMSSM and lattice QCD

Joel Giedt*,¹ Anthony W. Thomas†,² and Ross D. Young‡³

¹*Department of Physics, Applied Physics, and Astronomy,
Rensselaer Polytechnic Institute, 110 Eighth Street, Troy, New York 12180-3590 USA*

²*Jefferson Lab, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA
College of William and Mary, Williamsburg, Virginia 23187, USA*

³*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA*

Recent lattice measurements have given accurate estimates of the light and strange quark condensates in the proton. We use these new results to significantly improve the dark matter predictions in a set of benchmark models that represent different scenarios in the constrained minimal supersymmetric standard model (CMSSM). Because the predicted cross sections are at least an order of magnitude smaller than previously suggested, our results have significant consequences for dark matter searches.

PACS numbers: 12.60.Jv, 95.35.+d, 11.15.Ha, 11.30.Rd

The astronomical evidence for dark matter [1, 2] presents modern physics with some of its greatest challenges. We need to find ways to detect it directly and a number of very sophisticated searches are underway [3, 4, 5], with at times confusing results [6, 7]. To guide those searches we need theoretical models for what the dark matter might be and what cross section one might expect it to have for scattering from hadronic matter. The constrained minimal supersymmetric extensions of the Standard Model (CMSSM—see [8] for a review) have the advantage that they are consistent with modern nuclear and particle physics while incorporating the compelling concepts of supersymmetry, coupling unification, and viable cold dark matter [9]. These models have over time been tuned to ensure that they are consistent with the latest constraints on relic abundance [10, 11, 12, 13], with the favoured dark matter candidate being a neutralino with mass of order 100 GeV and a density of a few per litre.

Extensive studies of the spin-independent interaction of neutralinos with hadronic matter have established that the cross section is determined by their scalar coupling to the light and strange quark sigma commutators $((m_u + m_d)\langle\bar{u}u + \bar{d}d\rangle/2$ and $m_s\langle\bar{s}s\rangle$); see [12, 13] and references therein. Given that the favoured values of these quantities have been typically 50 MeV [14] and 300 MeV [15], respectively, the dark matter cross section has been dominated by the strange quark content of the proton. Realizing the importance of these quantities Ellis *et al.* recently made a plea for more accurate experimental data for them [12]. While this seems unlikely in the short term, at least, the answer to their plea has recently been provided from an unexpected source, through the study of octet baryon masses as a function of quark

mass. Indeed, a sophisticated chiral analysis of recent lattice measurements has yielded surprisingly accurate estimates of these sigma commutators [16]. Subsequent lattice computations have found consistent results [17].

In this Letter, we use the precise new values of the light and strange quark sigma commutators obtained in Refs. [16, 17] to update the cross sections predicted within the benchmark CMSSM models that have been studied for several years by Ellis *et al.* [12] and their collaborators [10, 11]. We show that the sizeable reduction in the strange sigma term from the values previously favoured leads to a rather dramatic reduction in the expected neutralino cross sections, with important implications for the interpretation of experiments to search for dark matter.

Sigma terms from lattice QCD. The sigma terms of the nucleon, for each quark flavour, are given by the scalar form factors evaluated at $t = 0$, denoted by

$$\sigma_q = m_q \langle N | \bar{q}q | N \rangle. \quad (1)$$

The nucleon scalar form factor is difficult to directly probe in experiment. The light-quark sigma term, $\sigma_\ell = (m_u + m_d)\langle\bar{u}u + \bar{d}d\rangle/2$, has most reliably been accessed by invoking a chiral low-energy relation between π - N scattering and the scalar form factor [14, 18, 19],

$$\Sigma_{\pi N} \equiv \sigma_\ell = \Sigma_{\pi N}^{\text{CD}} - \Delta_R - \Delta_\sigma. \quad (2)$$

The remainder term, Δ_R , describes a correction to the low-energy theorem and is estimated to be less than 2 MeV [19, 20]. The shift in the scalar form factor can be inferred from a dispersion analysis and found to be rather large [21], $\Delta_\sigma \equiv \sigma_\ell(2m_\pi^2) - \sigma_\ell = 15.4 \pm 0.4$ MeV. $\Sigma_{\pi N}^{\text{CD}}$ is the Born-subtracted, isoscalar πN scattering amplitude evaluated at the (unphysical) Cheng-Dashen point. An early experimental extraction [22] gave $\Sigma_{\pi N}^{\text{CD}} = 64 \pm 8$ MeV, to be compared with more a recent determination $\Sigma_{\pi N}^{\text{CD}} = 79 \pm 7$ MeV [23]. These two values lead to light-quark sigma terms of

$$\sigma_\ell = 45 \pm 8 \text{ MeV} \quad \text{and} \quad 64 \pm 7 \text{ MeV}, \quad (3)$$

*giedtj@rpi.edu
†awthomas@jlab.org
‡ryoung@anl.gov

respectively. These are to be compared with the recent lattice QCD determination $\sigma_\ell = 47 \pm 9 \text{ MeV}$ [16]. While this lattice analysis tends to favour the lower value, it is not inconsistent with the higher extraction. We use the lattice determination in the current analysis.

Extracting the strangeness sigma term is significantly more challenging, because the same prescription would lead to both a poorly converged low-energy relation and a large extrapolation of K - N scattering to the Cheng-Dashen point — see Reya [24]. A much more practical approach has been to resolve the patterns of $\text{SU}(3)$ breaking among the baryon octet [15, 25, 26]. In essence, the baryon masses give guidance with respect to the symmetry breaking component

$$\sigma_0 = m_\ell \langle N | \bar{u}u + \bar{d}d - 2\bar{s}s | N \rangle, \quad (4)$$

with $m_\ell = (m_u + m_d)/2$. The commonly reported value $\sigma_0 = 36 \pm 7 \text{ MeV}$ is based on the phenomenological analysis of corrections to the linear mass-splittings relations [25], with further uncertainties from yet higher order quantified by [26]. The difference between the estimated σ_0 and the extracted σ_ℓ then gives a best estimate for the strange-quark sigma term, as related by

$$\frac{\sigma_s}{\sigma_\ell} = \frac{m_s}{2m_\ell} \left(\frac{\Sigma_{\pi N} - \sigma_0}{\Sigma_{\pi N}} \right). \quad (5)$$

In contrast, lattice QCD allows one to directly study the quark mass dependence of the baryon masses from first principles. New lattice results, based on an $\text{SU}(3)$ baryon mass analysis [16] and a novel application of the Feynman-Hellman theorem at the correlator level [17], find $\sigma_s = 31 \pm 15 \text{ MeV}$ and $59 \pm 10 \text{ MeV}$, respectively. An important feature of these calculations is the demonstrated internal consistency, where both studies have successfully shown the ability to predict the mass-splittings between simulations with different strange quark mass parameters. We argue that these results show a vast improvement over the earlier phenomenological analyses. This, in turn, implies that it is appropriate to update the earlier estimates of dark matter cross sections. In the present analysis, we assume that the reported uncertainties in the lattice QCD determinations of the strange quark sigma term are independent and use a naïve average of the two results, $\sigma_s = 50 \pm 8 \text{ MeV}$. We report the results of our analysis at the 95% confidence level to ensure that the conclusions are minimally sensitive to our input.

Constrained MSSM. The constrained minimal supersymmetric standard model (CMSSM) is inspired by supergravity mediated scenarios of spontaneous supersymmetry breaking, in the context of supersymmetric grand unified theories [8]. Then one has a universal soft scalar mass m_0 , a universal gaugino mass $m_{1/2}$, a universal trilinear scalar coupling A_0 (which is set to zero in all the models considered here), and a higgsino mass μ . In prac-

tice one specifies the ratio of vacuum expectation values $\tan \beta = v_u/v_d$ of the Higgs fields H_u and H_d at the low energy scale (m_Z) and determines μ from this constraint; the sign of μ is also a parameter, and unification of gauge couplings is imposed. One advantage of the CMSSM class of models is its simplicity, allowing for detailed scans over parameter space. The benchmark models “A-M” of [10, 11] were selected to be representative of the regions of parameter space that yield neutralino dark matter with the correct abundance, consistent with WMAP constraints.

Because of the incredible accuracy of the WMAP results for relic cold dark matter abundance, a fine tuning of high scale parameters is typically required in order to have the lightest supersymmetric particle (LSP) fit this constraint. However, the dark matter cross section itself only changes by a few percent when this tuning is done. For instance, the tuning that we made (described below) in Model L, which is a 5% change to m_0 , only results in a 2% change in the spin-independent cross section. Thus the direct detection predictions are insensitive to the changes that may have to be made to bring the relic abundance into line with any future updates. Thus the cross section results presented here are robust in that sense.

Cross sections for benchmark models. We have computed the spin-independent cross section for neutralino dark matter for Benchmark Models A-M of [10, 11]. Three of the benchmark models were also studied in [12], where the reader may find the cross section formulae that we likewise use. While Ellis et al. use a private renormalization group evolution code, we have used the publicly available Softsusy [27] to compute the running parameters between the grand unified scale and the electroweak scale. (We did, however, compare our Softsusy results with those obtained using the code of Ellis et al. and found good agreement.) Minor modifications were necessary in two benchmark models, because of the slight differences between the renormalization group codes. These small shifts in high scale inputs were made in order to avoid a stau ($\tilde{\tau}$) LSP for models that are on the edge of the $\tilde{\tau}$ exclusion region of CMSSM (i.e., dark matter should be neutral). In Model J we changed m_0 from 285 GeV to 290 GeV. In Model L we changed m_0 from 300 to 315 GeV (305 and 310 were insufficient). As discussed above, these changes have an insignificant effect on the cross section values.

Our results for the spin independent cross section with the proton are shown in Figs. 1-3. In all cases we see that there is only a mild dependence on $\Sigma_{\pi N}$. Furthermore, most models have rather small cross sections, $\sigma_{SI} < 10^{-9} \text{ pb}$. We do not show Model L because it is essentially degenerate with Model I. Nor do we show Model D because it has such small cross sections ($\sigma_{SI} < 10^{-14} \text{ pb}$) that it is unobservable in the foreseeable future. We also computed the cross sections for the neutron and found them




FIG. 1: Proton, benchmark models (labeled by letters A, B,...), 95% CL using lattice inputs for sigma commutators. Remaining models are shown in other figures below, since they would overlap with these. The neutron predictions are not shown because they are basically degenerate with the proton cross sections on the logarithmic scale that is shown.

to be essentially degenerate with those of the proton. By way of contrast with previous estimates, in Fig. 4 we show σ_{SI} for Model C using σ_ℓ and σ_s extracted from lattice QCD with the determination of σ_s inferred from the phenomenological estimate for $\sigma_0 = 36$ MeV, using Eq. (5) above. Not only is the cross section calculated here, on the basis of the latest lattice QCD results, far more weakly dependent on $\Sigma_{\pi N}$, but it is also typically much smaller.

Details of quark flavor contributions. It is interesting to investigate the breakdown of how each quark flavor contributes to the overall cross section. First, we note that for the proton $\sigma_{SI} \propto |f_p|^2$, where

$$m_p f_p = \sum_{q=u,d,s} \frac{\alpha_{3q}}{m_q} f_{Tq}^p + \frac{2}{27} f_{TG}^p \sum_{q=c,b,t} \frac{\alpha_{3q}}{m_q},$$

$$f_{TG}^p = 1 - \sum_{q=u,d,s} f_{Tq}^p \quad (6)$$

The coefficients α_{3q} are determined by squark exchange and mixing coefficients of the LSP neutralino. The dimensionless sigma commutators $f_{Tq}^p = \sigma_q/m_p$ are taken from the lattice results, as explained earlier, whereas f_{TG}^p addresses the heavy flavor sigma commutators through SVZ relations, as discussed in the review [9]. We also find it interesting to compute the relative contribution of each quark flavor, defining $f_p = \sum_q f_q^p$. For instance, the bottom quark contribution $f_b^p = (2/27)m_p f_{TG}^p \alpha_{3b}/m_b$ turns out to be rather large. Its increased importance in our




FIG. 2: Cross section estimates (95% CL) for benchmark models C, I, J.




FIG. 3: Cross section estimates (95% CL) for benchmark models A, G.

computations can be traced to the fact that the lattice results give a smaller result for f_{Ts}^p , the dimensionless strange sigma commutator. This enhances f_{TG}^p , as can be seen from Eq. (6). Results are given in Table I for the point with maximum σ_{SI} , which occurs in Model L.

Conclusions. In this Letter we have shown that the recent lattice results [16, 17] for σ_ℓ and σ_s have a dramatic effect on predictions for direct detection of neutralino cold dark matter. Furthermore, the theoretical uncertainty is considerably smaller than in the traditional




FIG. 4: A comparison of our results (solid ellipse) for Model C, versus the traditional approach (dashed line) which relates the strange quark sigma commutator to the light quark one through Eq. (5) with $\sigma_0 = 36$ MeV.

Model	q	α_{3q}/m_q	f_{Tq}^p or f_{TG}^p	f_q^p/f_p
L (max σ_{SI}) $\sigma_{SI} =$ 2.8×10^{-9} pb	u	-1.019×10^{-09}	0.0280	0.0105
	d	-1.302×10^{-08}	"	0.1342
	c	-1.031×10^{-09}	0.8751	0.0261
	s	-1.522×10^{-08}	0.0689	0.3633
	t	-1.936×10^{-09}	0.8751	0.0462
	b	-1.670×10^{-08}	"	0.3984

TABLE I: Example breakdown of quark flavor contributions to σ_{SI} . This is for Model L at the maximum value of its cross section, within our 95% CL region.

approach, as was highlighted in Fig. 4. Unfortunately, the most important effect of the improved estimates is the reduction of the strange quark sigma commutator, which leads to rather small dark matter cross sections. On the other hand, even within the 95% confidence intervals, our estimates come with small errors; hence any observation of dark matter would be highly selective amongst the benchmark models that we have considered. In future work we hope to report on dark matter cross section predictions in models other than the CMSSM, such as the nonuniversal Higgs mass (NUHM) models, which are known [13] to allow for cross sections which are larger than in the CMSSM.

Acknowledgements. We thank Keith Olive for helpful discussions and for allowing us to use a version of his collaborations' RGE code in order to perform comparisons to the Softsusy output. JG was supported by

Rensselaer faculty development funds. This work was supported by the the U.S. Department of Energy contracts DE-AC05-06OR23177, under which Jefferson Science Associates, LLC operates Jefferson Laboratory and DE-AC02-06CH11357, under which UChicago Argonne, LLC, operates Argonne National Laboratory.

-
- [1] N. Jarosik *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **170**, 263 (2007) [arXiv:astro-ph/0603452].
 - [2] M. Tegmark *et al.* [SDSS Collaboration], *Phys. Rev. D* **69**, 103501 (2004) [arXiv:astro-ph/0310723].
 - [3] Z. Ahmed *et al.* [CDMS Collaboration], *Phys. Rev. Lett.* **102**, 011301 (2009) [arXiv:0802.3530 [astro-ph]].
 - [4] J. Angle *et al.*, *Phys. Rev. Lett.* **101**, 091301 (2008) [arXiv:0805.2939 [astro-ph]].
 - [5] C. E. Aalseth *et al.* [CoGeNT Collaboration], *Phys. Rev. Lett.* **101**, 251301 (2008) [Erratum-ibid. **102**, 109903 (2009)] [arXiv:0807.0879 [astro-ph]].
 - [6] C. Savage, K. Freese, P. Gondolo and D. Spolyar, arXiv:0901.2713 [astro-ph].
 - [7] R. Bernabei *et al.* [DAMA Collaboration], *Eur. Phys. J. C* **56**, 333 (2008) [arXiv:0804.2741 [astro-ph]].
 - [8] W. de Boer, *Prog. Part. Nucl. Phys.* **33** (1994) 201 [arXiv:hep-ph/9402266].
 - [9] G. Jungman, M. Kamionkowski and K. Griest, *Phys. Rept.* **267**, 195 (1996) [arXiv:hep-ph/9506380].
 - [10] M. Battaglia *et al.*, *Eur. Phys. J. C* **22** (2001) 535 [arXiv:hep-ph/0106204].
 - [11] M. Battaglia *et al.*, *Eur. Phys. J. C* **33** (2004) 273 [arXiv:hep-ph/0306219].
 - [12] J. R. Ellis, K. A. Olive and C. Savage, *Phys. Rev. D* **77**, 065026 (2008) [arXiv:0801.3656 [hep-ph]].
 - [13] J. Ellis, K. A. Olive and P. Sandick, arXiv:0905.0107 [hep-ph].
 - [14] J. Gasser, H. Leutwyler and M. E. Sainio, *Phys. Lett. B* **253**, 252 (1991).
 - [15] A. E. Nelson and D. B. Kaplan, *Phys. Lett. B* **192**, 193 (1987).
 - [16] R. D. Young and A. W. Thomas, arXiv:0901.3310 [hep-lat].
 - [17] D. Toussaint and W. Freeman [MILC Collaboration], arXiv:0905.2432 [hep-lat].
 - [18] T. P. Cheng and R. F. Dashen, *Phys. Rev. Lett.* **26**, 594 (1971).
 - [19] L. S. Brown, W. J. Pardee and R. D. Peccei, *Phys. Rev. D* **4**, 2801 (1971).
 - [20] V. Bernard, N. Kaiser and U. G. Meissner, *Phys. Lett. B* **389**, 144 (1996) [arXiv:hep-ph/9607245].
 - [21] J. Gasser, H. Leutwyler and M. E. Sainio, *Phys. Lett. B* **253**, 260 (1991).
 - [22] R. Koch, *Z. Phys. C* **15**, 161 (1982).
 - [23] M. M. Pavan, I. I. Strakovsky, R. L. Workman and R. A. Arndt, *PiN Newslett.* **16**, 110 (2002) [arXiv:hep-ph/0111066].
 - [24] E. Reya, *Rev. Mod. Phys.* **46**, 545 (1974).
 - [25] J. Gasser and H. Leutwyler, *Phys. Rept.* **87**, 77 (1982).
 - [26] B. Borasoy and U.-G. Meißner, *Annals Phys.* **254**, 192 (1997) [arXiv:hep-ph/9607432].
 - [27] B. C. Allanach, *Comput. Phys. Commun.* **143**, 305 (2002) [arXiv:hep-ph/0104145].